

# The block graph of a finite group

Alessandro Paolini

joint with J. Brough and Y. Liu

Technische Universität Kaiserslautern

Birmingham, August 7th 2017

$G$  finite group.  $p > 0$  prime,  $p \mid |G|$ .

**Goal:** relate the set  $\text{Irr}(G)$  of *ordinary* irreducible characters of  $G$  with the  $p$ -local structure of  $G$ .

$G$  finite group.  $p > 0$  prime,  $p \mid |G|$ .

**Goal:** relate the set  $\text{Irr}(G)$  of *ordinary* irreducible characters of  $G$  with the  $p$ -local structure of  $G$ .

Structure of the talk:

- 1) Setup
- 2) History
- 3) Block graphs
- 4) Results

- Let  $\mathcal{O}$  be the ring of algebraic integers in  $\mathbb{C}$ . If  $\chi \in \text{Irr}(G)$ , then

$$\omega_\chi(x) := \frac{\chi(x)}{\chi(1)} |x^G| \in \mathcal{O} \quad \text{for all } x \in G.$$

- Let  $\mathcal{O}$  be the ring of algebraic integers in  $\mathbb{C}$ . If  $\chi \in \text{Irr}(G)$ , then

$$\omega_\chi(x) := \frac{\chi(x)}{\chi(1)} |x^G| \in \mathcal{O} \quad \text{for all } x \in G.$$

- Let  $\mathfrak{m} \subseteq \mathcal{O}$  with  $p\mathcal{O} \subseteq \mathfrak{m}$ . Then  $\mathcal{O}/\mathfrak{m} \cong k$  with  $\text{char } k = p$ .

- Let  $\mathcal{O}$  be the ring of algebraic integers in  $\mathbb{C}$ . If  $\chi \in \text{Irr}(G)$ , then

$$\omega_\chi(x) := \frac{\chi(x)}{\chi(1)} |x^G| \in \mathcal{O} \quad \text{for all } x \in G.$$

- Let  $\mathfrak{m} \subseteq \mathcal{O}$  with  $p\mathcal{O} \subseteq \mathfrak{m}$ . Then  $\mathcal{O}/\mathfrak{m} \cong k$  with  $\text{char } k = p$ .
- Let  $\mathfrak{b}_i := e_i \mathcal{O}G$  for each primitive central idempotent  $e_i$  of  $\mathcal{O}G$ . Then

$$\mathcal{O}G := \mathfrak{b}_1 \oplus \cdots \oplus \mathfrak{b}_n$$

- Let  $\mathcal{O}$  be the ring of algebraic integers in  $\mathbb{C}$ . If  $\chi \in \text{Irr}(G)$ , then

$$\omega_\chi(x) := \frac{\chi(x)}{\chi(1)} |x^G| \in \mathcal{O} \quad \text{for all } x \in G.$$

- Let  $\mathfrak{m} \subseteq \mathcal{O}$  with  $p\mathcal{O} \subseteq \mathfrak{m}$ . Then  $\mathcal{O}/\mathfrak{m} \cong k$  with  $\text{char } k = p$ .
- Let  $\mathfrak{b}_i := e_i \mathcal{O}G$  for each primitive central idempotent  $e_i$  of  $\mathcal{O}G$ . Then

$$\mathcal{O}G := \mathfrak{b}_1 \oplus \cdots \oplus \mathfrak{b}_n \xrightarrow{\pi_{\mathfrak{m}}} kG = B_1 \oplus \cdots \oplus B_m,$$

and decompose  $kG$  into  $p$ -blocks of  $G$  by applying projection  $\pi_{\mathfrak{m}}$  over  $\mathfrak{m}$ .

- Let  $\mathcal{O}$  be the ring of algebraic integers in  $\mathbb{C}$ . If  $\chi \in \text{Irr}(G)$ , then

$$\omega_\chi(x) := \frac{\chi(x)}{\chi(1)} |x^G| \in \mathcal{O} \quad \text{for all } x \in G.$$

- Let  $\mathfrak{m} \subseteq \mathcal{O}$  with  $p\mathcal{O} \subseteq \mathfrak{m}$ . Then  $\mathcal{O}/\mathfrak{m} \cong k$  with  $\text{char } k = p$ .
- Let  $\mathfrak{b}_i := e_i \mathcal{O}G$  for each primitive central idempotent  $e_i$  of  $\mathcal{O}G$ . Then

$$\mathcal{O}G := \mathfrak{b}_1 \oplus \cdots \oplus \mathfrak{b}_n \xrightarrow{\pi_{\mathfrak{m}}} kG = B_1 \oplus \cdots \oplus B_m,$$

and decompose  $kG$  into  $p$ -blocks of  $G$  by applying projection  $\pi_{\mathfrak{m}}$  over  $\mathfrak{m}$ .

- This yields a partition of  $\text{Irr}(G)$  for each  $p$ . Let  $\chi, \psi \in \text{Irr}(G)$ .

$$\text{Irr}(G) = \bigsqcup_{B \text{ } p\text{-block of } G} \text{Irr}(B), \quad \chi \underset{p\text{-block}}{\overset{\text{same}}{\sim}} \psi \iff \pi_{\mathfrak{m}} \circ \omega_\chi = \pi_{\mathfrak{m}} \circ \omega_\psi.$$



- Let  $\mathcal{O}$  be the ring of algebraic integers in  $\mathbb{C}$ . If  $\chi \in \text{Irr}(G)$ , then

$$\omega_\chi(x) := \frac{\chi(x)}{\chi(1)} |x^G| \in \mathcal{O} \quad \text{for all } x \in G.$$

- Let  $\mathfrak{m} \subseteq \mathcal{O}$  with  $p\mathcal{O} \subseteq \mathfrak{m}$ . Then  $\mathcal{O}/\mathfrak{m} \cong k$  with  $\text{char } k = p$ .
- Let  $\mathfrak{b}_i := e_i \mathcal{O} G$  for each primitive central idempotent  $e_i$  of  $\mathcal{O} G$ . Then

$$\mathcal{O} G := \mathfrak{b}_1 \oplus \cdots \oplus \mathfrak{b}_n \xrightarrow{\pi_{\mathfrak{m}}} kG = B_1 \oplus \cdots \oplus B_m,$$

and decompose  $kG$  into  $p$ -blocks of  $G$  by applying projection  $\pi_{\mathfrak{m}}$  over  $\mathfrak{m}$ .

- This yields a partition of  $\text{Irr}(G)$  for each  $p$ . Let  $\chi, \psi \in \text{Irr}(G)$ .

$$\text{Irr}(G) = \bigsqcup_{B \text{ } p\text{-block of } G} \text{Irr}(B), \quad \chi \underset{p\text{-block}}{\overset{\text{same}}{\sim}} \psi \iff \pi_{\mathfrak{m}} \circ \omega_\chi = \pi_{\mathfrak{m}} \circ \omega_\psi.$$

- Finally, we define the *principal  $p$ -block*  $B_0(G)_p$  of  $G$  to be the unique  $p$ -block  $B$  of  $G$  such that  $\text{id}_G \in \text{Irr}(B)$ ,

$$\chi \in \text{Irr}(G), \quad \chi \in B_0(G)_p \iff \pi_{\mathfrak{m}} \circ \omega_\chi = \pi_{\mathfrak{m}} \circ \omega_{\text{id}_G}.$$

## Problem [Brauer '79]

Given a prime number  $p$ , find the relations between the properties of the  $p$ -blocks of characters of a finite group  $G$  and *structural* properties of  $G$ .

## Problem [Brauer '79]

Given a prime number  $p$ , find the relations between the properties of the  $p$ -blocks of characters of a finite group  $G$  and *structural* properties of  $G$ .

- If  $p \nmid |G|$  or  $\text{char } k = 0$ , get much structural global info from the character table, for instance:
  - obtain  $Z(G)$ , namely  $\{g \in G \mid |\chi(g)| = \chi(1) \text{ for every } \chi \in \text{Irr}(G)\}$
  - get every  $N \trianglelefteq G$  as  $\bigcap_{\chi \in \mathcal{I}} \ker(\chi)$ , for each  $\mathcal{I} \subseteq \text{Irr}(G)$ , where  $\ker(\chi) = \{g \in G \mid \chi(g) = \chi(1)\}$

## Problem [Brauer '79]

Given a prime number  $p$ , find the relations between the properties of the  $p$ -blocks of characters of a finite group  $G$  and *structural* properties of  $G$ .

- If  $p \nmid |G|$  or  $\text{char } k = 0$ , get much structural global info from the character table, for instance:
  - obtain  $Z(G)$ , namely  $\{g \in G \mid |\chi(g)| = \chi(1) \text{ for every } \chi \in \text{Irr}(G)\}$
  - get every  $N \trianglelefteq G$  as  $\bigcap_{\chi \in \mathcal{I}} \ker(\chi)$ , for each  $\mathcal{I} \subseteq \text{Irr}(G)$ , where  $\ker(\chi) = \{g \in G \mid \chi(g) = \chi(1)\}$
- We look at  $p$ -blocks when  $p \mid |G|$  for further local information .

## Proposition

Assume that there exists a  $p$ -block  $B$  such that  $B = \{\chi\}$ . Then  $O_p(G) = 1$ .

## Problem [Brauer '79]

Given a prime number  $p$ , find the relations between the properties of the  $p$ -blocks of characters of a finite group  $G$  and *structural* properties of  $G$ .

- If  $p \nmid |G|$  or  $\text{char } k = 0$ , get much structural global info from the character table, for instance:
  - obtain  $Z(G)$ , namely  $\{g \in G \mid |\chi(g)| = \chi(1) \text{ for every } \chi \in \text{Irr}(G)\}$
  - get every  $N \trianglelefteq G$  as  $\bigcap_{\chi \in \mathcal{I}} \ker(\chi)$ , for each  $\mathcal{I} \subseteq \text{Irr}(G)$ , where  $\ker(\chi) = \{g \in G \mid \chi(g) = \chi(1)\}$
- We look at  $p$ -blocks when  $p \mid |G|$  for further local information .

## Proposition

Assume that there exists a  $p$ -block  $B$  such that  $B = \{\chi\}$ . Then  $O_p(G) = 1$ .

## Theorem (Harris '85)

$B_0(G)_p$  is the only  $p$ -block if and only if

- $O_p(G) = F^*(G)$  if  $p \geq 3$ ; ( $F^*(G)$  is the generalized Fitting subgroup of  $G$ )
- $O_{2'}(G) = 1$  and all components of  $G$  are of type  $M_{22}$  and  $M_{24}$ , if  $p = 2$ .

- Compare now blocks at different primes  $p$  and  $r$ .

- Compare now blocks at different primes  $p$  and  $r$ .

**Theorem (Navarro, Willems '97)**

$p, r \mid |G|$ . Assume that there exist a  $p$ -block  $B_1$  and an  $r$ -block  $B_2$  such that  $\text{Irr}(B_1) = \text{Irr}(B_2)$ . If  $G$  is  $p$ -solvable or  $r$ -solvable, then  $\text{Irr}(B_1) = \{\chi\} = \text{Irr}(B_2)$ .

- Compare now blocks at different primes  $p$  and  $r$ .

### Theorem (Navarro, Willems '97)

$p, r \mid |G|$ . Assume that there exist a  $p$ -block  $B_1$  and an  $r$ -block  $B_2$  such that  $\text{Irr}(B_1) = \text{Irr}(B_2)$ . If  $G$  is  $p$ -solvable or  $r$ -solvable, then  $\text{Irr}(B_1) = \{\chi\} = \text{Irr}(B_2)$ .

### $p$ - or $r$ -solvability is necessary (Bessenrodt)

$G = 6.A_7$ . There exist a 5-block  $B_1$  and a 7-block  $B_2$ , with  $\text{Irr}(B_1) = \text{Irr}(B_2)$ , and  $|\text{Irr}(B_1)| = 5$ . The blocks  $B_1$  and  $B_2$  are *not* principal blocks.



- Compare now blocks at different primes  $p$  and  $r$ .

### Theorem (Navarro, Willems '97)

$p, r \mid |G|$ . Assume that there exist a  $p$ -block  $B_1$  and an  $r$ -block  $B_2$  such that  $\text{Irr}(B_1) = \text{Irr}(B_2)$ . If  $G$  is  $p$ -solvable or  $r$ -solvable, then  $\text{Irr}(B_1) = \{\chi\} = \text{Irr}(B_2)$ .

### $p$ - or $r$ -solvability is necessary (Bessenrodt)

$G = 6.A_7$ . There exist a 5-block  $B_1$  and a 7-block  $B_2$ , with  $\text{Irr}(B_1) = \text{Irr}(B_2)$ , and  $|\text{Irr}(B_1)| = 5$ . The blocks  $B_1$  and  $B_2$  are *not* principal blocks.

*Our focus now is on the study of principal blocks  $B_0(G)_p$ , for  $p \mid |G|$ .*

- Compare now blocks at different primes  $p$  and  $r$ .

### Theorem (Navarro, Willems '97)

$p, r \mid |G|$ . Assume that there exist a  $p$ -block  $B_1$  and an  $r$ -block  $B_2$  such that  $\text{Irr}(B_1) = \text{Irr}(B_2)$ . If  $G$  is  $p$ -solvable or  $r$ -solvable, then  $\text{Irr}(B_1) = \{\chi\} = \text{Irr}(B_2)$ .

### $p$ - or $r$ -solvability is necessary (Bessenrodt)

$G = 6.A_7$ . There exist a 5-block  $B_1$  and a 7-block  $B_2$ , with  $\text{Irr}(B_1) = \text{Irr}(B_2)$ , and  $|\text{Irr}(B_1)| = 5$ . The blocks  $B_1$  and  $B_2$  are *not* principal blocks.

*Our focus now is on the study of principal blocks  $B_0(G)_p$ , for  $p \mid |G|$ .*

### Theorem (Bessenrodt, Navarro, Olsson, Tiep '07)

Let  $p, r$  such that  $\text{Irr}(B_0(G)_p) = \text{Irr}(B_0(G)_r)$ . Then neither  $p$  nor  $r$  divide  $|G|$ . In particular,  $\text{Irr}(B_0(G)_p) = \{id_G\}$ .

- Compare now blocks at different primes  $p$  and  $r$ .

### Theorem (Navarro, Willems '97)

$p, r \mid |G|$ . Assume that there exist a  $p$ -block  $B_1$  and an  $r$ -block  $B_2$  such that  $\text{Irr}(B_1) = \text{Irr}(B_2)$ . If  $G$  is  $p$ -solvable or  $r$ -solvable, then  $\text{Irr}(B_1) = \{\chi\} = \text{Irr}(B_2)$ .

### $p$ - or $r$ -solvability is necessary (Bessenrodt)

$G = 6.A_7$ . There exist a 5-block  $B_1$  and a 7-block  $B_2$ , with  $\text{Irr}(B_1) = \text{Irr}(B_2)$ , and  $|\text{Irr}(B_1)| = 5$ . The blocks  $B_1$  and  $B_2$  are *not* principal blocks.

*Our focus now is on the study of principal blocks  $B_0(G)_p$ , for  $p \mid |G|$ .*

### Theorem (Bessenrodt, Navarro, Olsson, Tiep '07)

Let  $p, r$  such that  $\text{Irr}(B_0(G)_p) = \text{Irr}(B_0(G)_r)$ . Then neither  $p$  nor  $r$  divide  $|G|$ . In particular,  $\text{Irr}(B_0(G)_p) = \{id_G\}$ .

- Criteria for nilpotency, solvability, simplicity in terms of principal blocks related to different primes motivate the definition of *block graphs*.


**Main definition: block graph  $\Gamma_B(G)$**

**Main definition: block graph  $\Gamma_B(G)$** 

- vertices:  $\pi(G) = \{p \text{ prime} \mid p \text{ divides } |G|\}$ ;


## Main definition: block graph $\Gamma_B(G)$

- vertices:  $\pi(G) = \{p \text{ prime} \mid p \text{ divides } |G|\}$ ;
- $p \neq r$  are linked iff there exists a character  $\chi$  in  $\text{Irr}(B_0(G)_p) \cap \text{Irr}(B_0(G)_r)$  with  $\chi \neq \text{id}_G$ .



## Main definition: block graph $\Gamma_B(G)$

- vertices:  $\pi(G) = \{p \text{ prime} \mid p \text{ divides } |G|\}$ ;
- $p \neq r$  are linked iff there exists a character  $\chi$  in  $\text{Irr}(B_0(G)_p) \cap \text{Irr}(B_0(G)_r)$  with  $\chi \neq \text{id}_G$ .

$$1 \neq \chi \in \text{Irr}(B_0(G)_p) \cap \text{Irr}(B_0(G)_r)$$



## Theorem (Block graphs of nilpotent groups, Bessenrodt-Zhang '08)

Let  $|G| = p_1^{a_1} \cdots p_n^{a_n}$ . Then  $G$  is nilpotent if and only if  $\text{Irr}(B_0(G)_{p_i}) \cap \text{Irr}(B_0(G)_{p_j}) = 1$  for every  $i \neq j$ .

$$\begin{array}{cccc} \bullet & \bullet & \cdot & \bullet \\ p_1 & p_2 & & p_n \end{array}$$

## Main definition: block graph $\Gamma_B(G)$

- vertices:  $\pi(G) = \{p \text{ prime} \mid p \text{ divides } |G|\}$ ;
- $p \neq r$  are linked iff there exists a character  $\chi$  in  $\text{Irr}(B_0(G)_p) \cap \text{Irr}(B_0(G)_r)$  with  $\chi \neq \text{id}_G$ .


$$1 \neq \chi \in \text{Irr}(B_0(G)_p) \cap \text{Irr}(B_0(G)_r)$$


## Theorem (Block graphs of nilpotent groups, Bessenrodt-Zhang '08)

Let  $|G| = p_1^{a_1} \cdots p_n^{a_n}$ . Then  $G$  is nilpotent if and only if  $\text{Irr}(B_0(G)_{p_i}) \cap \text{Irr}(B_0(G)_{p_j}) = 1$  for every  $i \neq j$ .




- *Remark:* no universal bound on  $\text{diam}(\Gamma_B(G))$ . Take  $G = M_1 \times \cdots \times M_{n-1}$ , with each of the  $M_i$  non-nilpotent,  $|M_i| = p_i p_{i+1}$ .





## Main definition: block graph $\Gamma_B(G)$

- vertices:  $\pi(G) = \{p \text{ prime} \mid p \text{ divides } |G|\}$ ;
- $p \neq r$  are linked iff there exists a character  $\chi$  in  $\text{Irr}(B_0(G)_p) \cap \text{Irr}(B_0(G)_r)$  with  $\chi \neq \text{id}_G$ .


$$1 \neq \chi \in \text{Irr}(B_0(G)_p) \cap \text{Irr}(B_0(G)_r)$$


## Theorem (Block graphs of nilpotent groups, Bessenrodt-Zhang '08)

Let  $|G| = p_1^{a_1} \cdots p_n^{a_n}$ . Then  $G$  is nilpotent if and only if  $\text{Irr}(B_0(G)_{p_i}) \cap \text{Irr}(B_0(G)_{p_j}) = 1$  for every  $i \neq j$ .



- *Remark:* no universal bound on  $\text{diam}(\Gamma_B(G))$ . Take  $G = M_1 \times \cdots \times M_{n-1}$ , with each of the  $M_i$  non-nilpotent,  $|M_i| = p_i p_{i+1}$ .




## Theorem ( $\Gamma_B(S)$ for $S$ alternating or sporadic, Bessenrodt-Zhang '08)

- (i) If  $n \geq 4$ , then  $\Gamma_B(\text{Alt}(n))$  is a complete graph.



## Main definition: block graph $\Gamma_B(G)$

- vertices:  $\pi(G) = \{p \text{ prime} \mid p \text{ divides } |G|\}$ ;
- $p \neq r$  are linked iff there exists a character  $\chi$  in  $\text{Irr}(B_0(G)_p) \cap \text{Irr}(B_0(G)_r)$  with  $\chi \neq \text{id}_G$ .




## Theorem (Block graphs of nilpotent groups, Bessenrodt-Zhang '08)

Let  $|G| = p_1^{a_1} \cdots p_n^{a_n}$ . Then  $G$  is nilpotent if and only if  $\text{Irr}(B_0(G)_{p_i}) \cap \text{Irr}(B_0(G)_{p_j}) = 1$  for every  $i \neq j$ .





- *Remark:* no universal bound on  $\text{diam}(\Gamma_B(G))$ . Take  $G = M_1 \times \cdots \times M_{n-1}$ , with each of the  $M_i$  non-nilpotent,  $|M_i| = p_i p_{i+1}$ .



## Theorem ( $\Gamma_B(S)$ for $S$ alternating or sporadic, Bessenrodt-Zhang '08)

- If  $n \geq 4$ , then  $\Gamma_B(\text{Alt}(n))$  is a complete graph.
- $S$  sporadic. Then  $\Gamma_B(S)$  is complete iff  $S \neq J_1$   $[(p, r) = (3, 5)]$  and  $S \neq J_4$ ,  $[(p, r) = (5, 7)]$ .



- **Question:** can we complete the determination of block graphs of finite simple groups?

- **Question:** can we complete the determination of block graphs of finite simple groups? YES!

- **Question:** can we complete the determination of block graphs of finite simple groups? YES!

### Theorem (Brough, Liu, P. '17)

*Let  $S$  be a finite simple group of Lie type. Then  $\Gamma_B(S)$  is complete.*

- **Question:** can we complete the determination of block graphs of finite simple groups? YES!

### Theorem (Brough, Liu, P. '17)

*Let  $S$  be a finite simple group of Lie type. Then  $\Gamma_B(S)$  is complete.*

*Idea of proof.* Fixed  $\ell_1, \ell_2 \mid |G|$ , need to find  $1 \neq \chi \in \text{Irr}(B_0(S))_{\ell_1} \cap \text{Irr}(B_0(S))_{\ell_2}$ .

- **Question:** can we complete the determination of block graphs of finite simple groups? YES!

### Theorem (Brough, Liu, P. '17)

*Let  $S$  be a finite simple group of Lie type. Then  $\Gamma_B(S)$  is complete.*

*Idea of proof.* Fixed  $\ell_1, \ell_2 \mid |G|$ , need to find  $1 \neq \chi \in \text{Irr}(B_0(S))_{\ell_1} \cap \text{Irr}(B_0(S))_{\ell_2}$ .

- Let  $\mathbb{G}$  be the algebraic group with Frobenius endomorphism  $F$  over  $\mathbb{F}_q$  associated to  $S$ . If  $F$  is very twisted, the claim follows from [Hiss '10].

- **Question:** can we complete the determination of block graphs of finite simple groups? YES!

### Theorem (Brough, Liu, P. '17)

Let  $S$  be a finite simple group of Lie type. Then  $\Gamma_B(S)$  is complete.

*Idea of proof.* Fixed  $\ell_1, \ell_2 \mid |G|$ , need to find  $1 \neq \chi \in \text{Irr}(B_0(S))_{\ell_1} \cap \text{Irr}(B_0(S))_{\ell_2}$ .

- Let  $\mathbb{G}$  be the algebraic group with Frobenius endomorphism  $F$  over  $\mathbb{F}_q$  associated to  $S$ . If  $F$  is very twisted, the claim follows from [Hiss '10].
- Let  $F$  be not very twisted. Define

$$e_j := \text{multiplicative order of } q \begin{cases} \text{modulo } \ell_j \text{ if } \ell_j \text{ is odd,} \\ \text{modulo } 4 \text{ if } \ell_j = 2. \end{cases}$$



- **Question:** can we complete the determination of block graphs of finite simple groups? YES!

### Theorem (Brough, Liu, P. '17)

Let  $S$  be a finite simple group of Lie type. Then  $\Gamma_B(S)$  is complete.

*Idea of proof.* Fixed  $\ell_1, \ell_2 \mid |G|$ , need to find  $1 \neq \chi \in \text{Irr}(B_0(S))_{\ell_1} \cap \text{Irr}(B_0(S))_{\ell_2}$ .

- Let  $\mathbb{G}$  be the algebraic group with Frobenius endomorphism  $F$  over  $\mathbb{F}_q$  associated to  $S$ . If  $F$  is very twisted, the claim follows from [Hiss '10].
- Let  $F$  be not very twisted. Define

$$e_i := \text{multiplicative order of } q \begin{cases} \text{modulo } \ell_i \text{ if } \ell_i \text{ is odd,} \\ \text{modulo } 4 \text{ if } \ell_i = 2. \end{cases}$$

- We construct Levi subgroups  $\mathbb{L}_i$  of  $\mathbb{G}$ ,  $i = 1, 2$ , centralizers of *Sylow*  $e_i$ -tori. Let  $R_{\mathbb{L}_i}^{\mathbb{G}}(1_{\mathbb{L}_i^F})$  be the induced (virtual) *Lusztig character*.

- **Question:** can we complete the determination of block graphs of finite simple groups? YES!

### Theorem (Brough, Liu, P. '17)

Let  $S$  be a finite simple group of Lie type. Then  $\Gamma_B(S)$  is complete.

*Idea of proof.* Fixed  $\ell_1, \ell_2 \mid |G|$ , need to find  $1 \neq \chi \in \text{Irr}(B_0(S))_{\ell_1} \cap \text{Irr}(B_0(S))_{\ell_2}$ .

- Let  $\mathbb{G}$  be the algebraic group with Frobenius endomorphism  $F$  over  $\mathbb{F}_q$  associated to  $S$ . If  $F$  is very twisted, the claim follows from [Hiss '10].
- Let  $F$  be not very twisted. Define

$$e_i := \text{multiplicative order of } q \begin{cases} \text{modulo } \ell_i \text{ if } \ell_i \text{ is odd,} \\ \text{modulo } 4 \text{ if } \ell_i = 2. \end{cases}$$

- We construct Levi subgroups  $\mathbb{L}_i$  of  $\mathbb{G}$ ,  $i = 1, 2$ , centralizers of Sylow  $e_i$ -tori. Let  $R_{\mathbb{L}_i}^{\mathbb{G}}(1_{\mathbb{L}_i^F})$  be the induced (virtual) Lusztig character. We use:

### Theorem (Kessar, Malle '15)

The constituents of  $R_{\mathbb{L}_i}^{\mathbb{G}}(1_{\mathbb{L}_i^F})$  lie inside  $B_0(G)_{\ell_i}$ ,  $i = 1, 2$ .

- **Question:** can we complete the determination of block graphs of finite simple groups? YES!

### Theorem (Brough, Liu, P. '17)

Let  $S$  be a finite simple group of Lie type. Then  $\Gamma_B(S)$  is complete.

*Idea of proof.* Fixed  $\ell_1, \ell_2 \mid |G|$ , need to find  $1 \neq \chi \in \text{Irr}(B_0(S))_{\ell_1} \cap \text{Irr}(B_0(S))_{\ell_2}$ .

- Let  $\mathbb{G}$  be the algebraic group with Frobenius endomorphism  $F$  over  $\mathbb{F}_q$  associated to  $S$ . If  $F$  is very twisted, the claim follows from [Hiss '10].
- Let  $F$  be not very twisted. Define

$$e_i := \text{multiplicative order of } q \begin{cases} \text{modulo } \ell_i \text{ if } \ell_i \text{ is odd,} \\ \text{modulo } 4 \text{ if } \ell_i = 2. \end{cases}$$

- We construct Levi subgroups  $\mathbb{L}_i$  of  $\mathbb{G}$ ,  $i = 1, 2$ , centralizers of Sylow  $e_i$ -tori. Let  $R_{\mathbb{L}_i}^{\mathbb{G}}(1_{\mathbb{L}_i^F})$  be the induced (virtual) Lusztig character. We use:

### Theorem (Kessar, Malle '15)

The constituents of  $R_{\mathbb{L}_i}^{\mathbb{G}}(1_{\mathbb{L}_i^F})$  lie inside  $B_0(G)_{\ell_i}$ ,  $i = 1, 2$ .

- The claim follows from  $\langle R_{\mathbb{L}_1}^{\mathbb{G}}(1_{\mathbb{L}_1^F}), R_{\mathbb{L}_2}^{\mathbb{G}}(1_{\mathbb{L}_2^F}) \rangle = 0$  and  $\langle R_{\mathbb{L}_i}^{\mathbb{G}}(1_{\mathbb{L}_i^F}), 1_{\mathbb{G}^F} \rangle = 1$ .

- We would like to detect the solvability of  $G$  via its block graph.

- We would like to detect the solvability of  $G$  via its block graph.

### Theorem (Brough, Liu, P. '17)

*Let  $G$  be a finite group,  $p \mid |G|$ . If  $\Gamma_B(G)$  does not contain triangles with vertex  $p$ , then  $G$  is  $p$ -solvable. In particular (by the Feit–Thompson theorem) if  $\Gamma_B(G)$  does not contain triangles with vertex 2, then  $G$  is solvable.*

- We would like to detect the solvability of  $G$  via its block graph.

### Theorem (Brough, Liu, P. '17)

*Let  $G$  be a finite group,  $p \mid |G|$ . If  $\Gamma_B(G)$  does not contain triangles with vertex  $p$ , then  $G$  is  $p$ -solvable. In particular (by the Feit–Thompson theorem) if  $\Gamma_B(G)$  does not contain triangles with vertex 2, then  $G$  is solvable.*

**The viceversa does not hold.**  $G = C_5^3 \rtimes \text{Sym}(3)$

Bessenrodt–Zhang '11: if  $\pi(G) = \pi_1 \sqcup \pi_2$ , then  $\pi_1$  and  $\pi_2$  are disconnected in  $\Gamma_B(G)$  if and only if  $G = O_{\pi_1}(G) \times O_{\pi_2}(G)$ .

- We would like to detect the solvability of  $G$  via its block graph.

### Theorem (Brough, Liu, P. '17)

Let  $G$  be a finite group,  $p \mid |G|$ . If  $\Gamma_B(G)$  does not contain triangles with vertex  $p$ , then  $G$  is  $p$ -solvable. In particular (by the Feit–Thompson theorem) if  $\Gamma_B(G)$  does not contain triangles with vertex 2, then  $G$  is solvable.

### The viceversa does not hold. $G = C_5^3 \rtimes \text{Sym}(3)$

Bessenrodt–Zhang '11: if  $\pi(G) = \pi_1 \sqcup \pi_2$ , then  $\pi_1$  and  $\pi_2$  are disconnected in  $\Gamma_B(G)$  if and only if  $G = O_{\pi_1}(G) \times O_{\pi_2}(G)$ .

*Idea of proof.* Reduction argument.

- If  $G$  is minimal not  $p$ -solvable, there is a minimal normal subgroup of  $G$  isomorphic to  $S^t$  with  $S$  nonabelian simple,  $p \mid |S|$ .
- Let  $A := \text{Aut}(S)$  and  $M := A^t \cap G$ . Then  $C_M(S) \leq N_M(S) = M$ , and  $\overline{M} := M/C_M(S)$  is almost simple with  $\text{Soc}(\overline{M}) = S$ , that is,  $S \leq \overline{M} \leq A$ .
- $S$  not of Lie type: [Bessenrodt–Zhang '08].  $S$  of Lie type: via *Zsigmondy primes*, find  $r, \ell \in \pi(S)$  such that  $\{p, r, \ell\}$  is a triangle in  $\Gamma_B(G)$ .